

---

## HAAO Antibody

**Subcategory:** Rabbit Polyclonal Antibody

**Cat. No.:** 254102

**Unit:** 0.1 mg

### Description:

HAAO (3-Hydroxyanthranilate 3,4-dioxygenase) is a monomeric cytosolic protein of the family of intramolecular dioxygenases containing non-heme ferrous iron. It is widely distributed in peripheral organs, such as liver and kidney, and is present in low amounts in the central nervous system. This enzyme participates in tryptophan metabolism. It employs one cofactor, iron. HAAO catalyzes the synthesis of quinolinic acid (QUIN) from 3-hydroxyanthranilic acid. QUIN is an excitotoxin whose toxicity is mediated by its ability to activate glutamate N-methyl-D-aspartate receptors. Increased cerebral levels of QUIN may participate in the pathogenesis of neurological and inflammatory disorders. HAAO has been suggested to play a role in disorders associated with altered tissue levels of QUIN. Furthermore, recent study shows that HAAO are excellent candidate biomarkers for detecting ovarian cancer.

**Isotype:** Rabbit Ig

**Applications:** E, WB, IHC

**Species Reactivity:** H, M, R

**Format:** Each vial contains 0.1 ml IgG in PBS pH 7.4 with 0.02% sodium azide. Antibody was purified by immunogen affinity chromatography.

**Alternate Names:** HAAO; 3-hydroxyanthranilate 3;4-dioxygenase; 3-hydroxyanthranilate oxygenase; 3-HAO; HAD; HAO

**Accession No.:** NP\_036337

**Antigen:** KLH-conjugated synthetic peptide encompassing a sequence within human HAAO.

**Application Notes:** E: 1:500-1:1,000; WB: 1:100-1:500; IHC: 1:00-1:500

**Storage:** Store at -20°C. Minimize freeze-thaw cycles.

Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.