

## From Biology to Discovery™

## **HGFR Antibody**

Subcategory: Rabbit Polyclonal Antibody

Cat. No.: 200113 Unit: 0.1 mg

## Description:

HGFR (MET), a tyrosine protein kinase, is the receptor for the hepatocyte growth factor and scater factor. Activation of HGFR after rearrangement with the TPR gene produces an oncogenic protein c-MET. Defects in HGFR/MET are a cause of hepatocellular carcinoma and renal carcinoma.

**Isotype:** Rabbit Ig **Applications:** E, WB

Species Reactivity: H, M, R, Mk, P, Rb, G

**Format:** Each vial contains 0.1 mg lgG in 0.1 ml (1 mg/ml) of PBS pH7.4 with 0.09% sodium azide. Antibody was purified

by Protein-G affinity chromatography.

**Alternate Names:** Hepatocyte growth factor receptor; HGF receptor; Scatter factor receptor; SF receptor; HGF/SF receptor; Met proto-oncogene tyrosine kinase; c-Met

Accession No.: P08581

Antigen: KLH-conjugated synthetic peptide encompassing a

sequence within the N-term region of human HGFR. **Application Notes:** E: 1:500-1:1,000; WB: 1:100-1:500

**Storage:** Store at -20°C. Minimize freeze-thaw cycles. Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.