
IDH2 Antibody

Subcategory: Rabbit Polyclonal Antibody

Cat. No.: 254307

Unit: 0.1 mg

Description:

Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Two NADP(+)-dependent isocitrate dehydrogenases have been found as homodimer: IDH1 is predominantly cytosolic and peroxisomal and IDH2 is mitochondrial. Both IDH1 and IDH2 play significant roles in the tricarboxylic acid cycle (TCA), and defects in IDH1 as well as IDH2 have been implicated in the development of glioma as well as other malignancies.

Isotype: Rabbit Ig

Applications: E, WB, IHC

Species Reactivity: H, M, R

Format: Each vial contains 0.1 ml IgG in PBS pH 7.4 with 0.02% sodium azide. Antibody was purified by immunogen affinity chromatography.

Alternate Names: IDH2; Mitochondrial NADP-isocitrate dehydrogenase; IDH; IDP; IDHM; IDPM; ICD-M; mNADP-IDH

Accession No.: NP_002159

Antigen: KLH-conjugated synthetic peptide encompassing a sequence within human IDH2.

Application Notes: E: 1:500-1:1,000; WB: 1:100-1:500; IHC: 1:00-1:500

Storage: Store at -20°C. Minimize freeze-thaw cycles.

Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.